

EFFECTS OF THE DOCUMENTED LAND USE CHANGES ON THE ALBEDO AND EVAPOTRANSPIRATION IN EAST-HUNGARY

János Mika¹ Szilvia Horváth² and László Makra²

¹Hungarian Meteorological Service; P. O. Box 39, H-1675 Budapest, Hungary,
e-mail: mika.j@met.hu

²University of Szeged, Department of Climatology & Landscape Ecology,
P. O. Box 653, H-6701 Szeged, Hungary;
e-mail: lupu@geo.u-szeged.hu; makra@geo.u-szeged.hu

Abstract. Agricultural land use series are investigated in a plain catchment area of the Tisza River, almost identically represented by six administrative counties. Each county, commonly covering 34,000 km², is characterised by 70-80 % of managed vegetation. Effects of area coverage variations between the different plants are computed for the period 1951-2000 by applying results of a literature-based syntheses, specified for Hungary. These studies estimate surface-albedo values and proportion between real and potential evapotranspiration for the great majority of the plants grown in the region. Product of the plant-specific characteristics and the relative area coverage result in monthly series of surface albedo and real evapotranspiration. These values are related to the energy balance of the surface-atmosphere system by using a radiative-convective model adjusted for the given location. Two questions are investigated and positively answered: i) Are there monotonous trends in the given terms of the energy and water budget? ii) Are these changes comparable to the effects caused by other external forcings?

Key words: land-use, albedo, evapotranspiration, climate change, biofeedback

DIE WIRKUNG DER DOKUMENTIERTEN LANDNUTZUNG AUF DIE ALBEDO UND DIE EVAPOTRANSPIRATION IN OSTUNGARN

Zusammenfassung. In unserer Arbeit haben wir die landwirtschaftlichen Landnutzungsdaten auf dem Einzugsgebiet der Theiß (34 000 Km²) untersucht, das sich auf der großen ungarischen Tiefebene befindet. 70-80 % des Untersuchungsgebietes ist angebaut. Die Wirkungen der Gebiets-veränderungen der Pflanzenbedeckung sind mit Hilfe der ungarischen Fachliteratur zwischen 1951-2000 berechnet worden. Diese Studien geben eine Schätzung auf die Werte der Oberflächen-albedo, die Verhältnisse der wirklichen und möglichen Evapotranspiration in Anbetracht auf die Kulturpflanzen auf dem Einzugsgebiet. Danach haben wir die Werte der Oberflächenalbedo, der Systemalbedo und der wirklichen Evapotranspiration mit Hilfe von den vorgenannten meteorologischen Daten und den Verhältnissen der Pflanzenbedeckung berechnet. Weiterhin haben wir festgestellt, dass diese Werte zu den Ergebnissen der Energiebilanz des Oberfläche-Atmosphäre Systems in Beziehung stehen, die mit Hilfe von dem Strahlungs und konvektiven Modell auf unserem Untersuchungsgebiet berechnet worden. In unserer Studie suchen wir die Antwort auf drei Fragen: i) Ob wir Monotontrende in der untersuchten Periode der Energie- und Wasserbilanz finden können? ii) Ob diese Änderungen mit die Wirkungen vergleichbar und positive beantwortet sind, die durch die anderen äußeren Zwängen verursacht sind?

Schlüsselworte: Landnutzung, Albedo, Evapotranspiration, Klimaänderung, Rückkopplung

1. Introduction

Sunshine radiation absorbed by the Earth-atmosphere system is about 240 Wm⁻². The greenhouse effect is expected to add about 6 Wm⁻² by the middle of the 21st century (IPCC, 1996). Consequently, recent and future climate change forcing considerations are based on calculations originating from 1 % change of the energy balance. This is a very little number compared either to measuring accuracy of most environmental physical parameters, or to relative error of model calculations. Hence, it is worth studying various external factors and feedback mechanisms which might influence climate (CLIVAR, 1998).

One set of the possible feedbacks can be focused by vegetation. Role of plant cover has already been mentioned in the literature since the 1970's, when two possible ways of surface modification, namely overgrazing in the subtropics and devastation of rain forests were counted to be potential causes of the global climate change (*Charney et al., 1977; Sagan et al, 1979*). Both changes express their effect on climate through the light-reflecting capacity of the surface, i.e. its *albedo*.

A likely biological feedback was found in two steps for Hungary. *Mika (1988)* established statistically that during the last 100 years, the 0.5 K hemispheric warming had been accompanied with 0.5-0.8 K temperature increase, 7-14 percent decrease of precipitation, 20 % increase of sunshine duration in the summer half-year. These changes correspond to ca. 60 % increase in frequency of dry months (i.e. soil moisture content < 30 % of field capacity) and to 8-10 % increase of global radiation at the surface.

In relation to these local changes, three possible feedback mechanisms, connected to surface albedo modifications, were quantified. These are the longer vegetation period, the less precipitation and the adequate alterations in the managed vegetation, all induce increase of the surface albedo. Characteristic albedo values are taken from an independent climatological synthesis of local albedo measurements (*Dávid, 1985*). In a study (*Mika et al., 1992*), the sum of these feedbacks was assessed to be -0.7 Wm^{-2} presuming changes in vegetation cover due to regional consequences of 0.5 K global warming in Hungary. This value is comparable to the radiative forcing of 100 ppm CO_2 increase.

The *present study* is aimed to analyse the effects of registered changes in the managed vegetation on evapotranspiration and surface albedo. *Sect. 2* describes the methods of computation for evapotranspiration, surface albedo and short-wave radiation balance of the earth-atmosphere system perturbed by changes of vegetation. *Sect. 3* introduces the investigated region in East-Hungary and the experienced changes of vegetation between 1951 and 2000. *Sect. 4* shows the results of computed changes in the physical parameters described in *Sect. 2*, also those compared to changes hypothetically caused by other factors. Limitations of these computations, in respect to general comparison of these regional processes to the regional effects of the global ones, are discussed in *Sect. 5*.

2. Estimation of evapotranspiration and albedo

2.1 Evapotranspiration

Evapotranspiration of natural surfaces cannot be measured directly; it should be assessed by calculation, first step of which is to determine the potential evapotranspiration, ET_p . The real evapotranspiration is influenced also by available water capacity of the soil and by transpiration capacity of plants. In our computations, the former effect is expressed by the *relative soil moisture*, w , which is the ratio of actual soil moisture content and field capacity, i.e. the water of the upper soil layers, available for plants.

Transpiration by plants is determined by the growing season and individual state of plants, which is considered by the *biological constant*, B , which is generally less than 1. *Table 1.* shows monthly values of the biological constant for some frequent agricultural plants (*Antal, 1968*). These factors of evapotranspiration, from a plant-covered surface, ET , are connected by the formula (1) (*Antal, 1968*), where w is the relative soil moisture, B the biological constant and ET_p the potential evapotranspiration (mm).

$$ET = \frac{w + B}{1 + B} \cdot w \cdot ET_p, \quad (1)$$

Our calculations of evapotranspiration are performed by using five-days averages, applying biological constants determined by *Posza and Stollár (1983)* similar to that followed by *Antal (1968)*. The B plant constants were evaluated for 19 species by field experiments performed at the Agrometeorological Observatories since the late sixties in Hungary, using Thornthwaite type compensation lysimeter (*Posza and Stollár, 1983*).

Table 1. Monthly average values of the *B* biological constant for the most frequent agricultural plants in Hungary (after *Antal, 1968*)

	April	May	June	July	Aug.	Sept.	Oct.
<i>maize</i>	0.00	0.06	0.31	0.89	0.69	0.17	0.00
<i>potato</i>	0.02	0.17	0.74	0.90	0.48	0.08	0.00
<i>sugar beet</i>	0.03	0.16	0.61	0.94	0.59	0.24	0.06
<i>alfalfa</i>	0.32	0.60	0.74	0.86	0.80	0.65	0.33
<i>red clover</i>	0.48	0.67	0.87	0.83	0.88	0.65	0.29
<i>wheat</i>	0.38	0.87	0.92	0.28			
<i>barley</i>	0.31	0.62	0.93	0.50			
<i>oat</i>	0.04	0.38	0.86	0.47			
<i>pea</i>	0.10	0.54	0.48				

For *w* the soil moisture data of *Dunkel (1994)* are used, which are based on monthly surface water balance computations, considering actual precipitation, temperature and vapour pressure. For the calculation of potential evaporation, the empirical regression (2) was used (*Antal, 1968*), where *a*, *b* *c* and *A* are empirical constants, *e_s* saturation water vapour pressure, *r* relative humidity, *t* mean air temperature.

$$ET_p = a[e_s(1-r)]^b \cdot (1 + A \cdot t)^c \quad (2)$$

In the followings, sums of evapotranspiration are calculated from long-term of climatic averages of *ET_p* and *w* in order to isolate effects of land-use changes on evapotranspiration. Historical variations of these values, driven by the meteorological anomalies, are introduced only for comparison in Sect. 4.1.

2.2 Surface albedo

Surface albedo is influenced by the type and state of soils, species of plant cover and its growing phase. *Dávid (1985)* synthesised surface albedo values of special literature, established according to plants and growing phases. Territory of Hungary can be divided to some plant-specific regions, according to the temporal differences of growing phases. *Dávid* established climatically specific surface albedos for these regions and groups of plants in ten days' resolution. Average monthly surface albedo of plants, most frequently raised in Hungary, are listed in *Table 2*. In *Table 2.*, Region 1 indicates southern part of East-Hungary, while Region 2 relates to its northern part, differing in average thermal and hydrological characteristics.

2.3 Radiation balance of the earth-atmosphere system

Change of energy balance, caused by the land use, was determined not only at the surface but to the surface-atmosphere system, as well. This makes possible to compare energy changes due to surface modification with primary energetic effects of CO₂-concentration or aerosols, estimation of which is more reliable at the top of the atmosphere.

For this aim, results of a former calculation (*Mika et al., 1993*), made by a radiative-convective model (*Práger and Kovács, 1988*), adapted from the Main Geophysical Observatory, St. Petersburg (*Karol and Frolkis, 1984*), was used, by freezing its convective adjustment and other feedback mechanisms to simulate primary changes of the radiative heating. The model is horizontally averaged with 16 levels of computation from 1000 hPa at the surface to 0.64 hPa at ca. 60 km from the surface. Step between the levels is 100 hPa in the troposphere. Internal parameters and astronomical conditions of the model are set for Budapest (47°26' N, 19°17' E). Cloud amount of different layers are from *Warren et al. (1985)*, adjusted to local climatology. Low- and medium-level clouds are considered as blackbodies for long-wave radiation. High-level cloudiness is simulated by 0.5 emissivity coefficient. Aerosol optical profile is adapted from *WMO (1983)* considering "continental background" aerosol.

Table 2. Average surface albedo of some plants (after Dávid, 1985)

Region	Surface albedo, %						
	Apr.	May	June	July	Aug.	Sept.	Oct.
	Wheat						
1.	18	20	33	23	21		
2.	18	20	21	23	21		
	Barley						
1.	17	20	21	23	21		
2.	17	19	21	23	21		
	Rye						
1.	18	20	21	23	20		
2.	18	19	21	23	20		
	Maize						
1.		15	18	23	23	24	25
2.		15	17	20	23	23	25
	Alfalfa						
1.&2.	23	20	23	22	23	23	19
	Potato						
1.	15	19	24	20	18	19	
2.	15	18	22	23	19	19	
	sugar beet						
1.	14	15	19	19	21	22	22
2.		14	18	19	20	22	22
	lawn (meadow, pasture lands)						
1.&2.	17	19	20	20	19	20	19
	forests (in leaf and conifers)						
1.&2.	14						

By this model we determined how surface albedo changes affect the short-wave energy balance of the surface-atmosphere system, R_s . Connection to the system albedo, α_s is:

$$\Delta R_s = -G_0 \cdot \Delta \alpha_s, \quad (3)$$

where G_0 is sunshine radiation at the top of the atmosphere. Dependence of the system albedo on α surface albedo is, with high accuracy, linear (Mika *et al.*, 1992) from which:

$$\Delta \alpha_s = k \cdot \Delta \alpha. \quad (4)$$

Unit change of surface albedo involves $k = 0.40 - 0.45$ change of system albedo in the examined seven months. Damped changes appearing at the outer boundary of the atmosphere can mainly be explained by the existence of cloudiness and limited transparency of the atmosphere. The higher coefficients characterise the summer period, when cloudiness is less and the optical length is shorter.

3. Land use in the East-Hungarian region

3.1 The selected region

The region, selected for the investigation, is the Hungarian plain-water catchment area of the Tisza River (Fig. 1), for which there was previously made a regional energy- and water balance model, as well (Mika *et al.*, 1991; 1998). In the followings, this region will be represented by six administrative counties, called Borsod-Abaúj-Zemplén, Szabolcs-Szatmár-Bereg, Hajdú-Bihar, Jász-Nagykun-Szolnok, Békés and Csongrád, considering the fact that land use data are officially published county by county. Natural vegetation of the Hungarian Plain represents the westernmost extension of this forest-steppe zone in Europe. This large landscape has always been characterised by high proportionality of managed vegetation. Recently, 74 % of the total administrative area (34,000 km²) is cultivated.




Fig. 1. The Tisza River sub-catchment in East-Hungary and the administrative counties, six of which approximately cover this region (34,000 km²)




Fig. 2. Total sown area of the plant groups in the six counties 1951-2000, hectare (1 ha = 0.01 km²)

3.2 Land use series

In this chapter it is briefly shown, how sown area of the different plants developed in six examined counties according to the data in the annual reference books of the *Central Statistical Office* (1951-2000), and from *Historical Statistical Contributions* (1971-79). The relative share of various plants was determined for the period 1951-2000.

The plants are also arranged into five groups, for easier interpretation, but original plants are used in the computations: *cereals* - wheat, rye, barley and rice; *fodder-plants* - maize, alfalfa, red clover, maize for silage, oat and cattle-turnip; *food and industrial plants* - sugar beet, tobacco, sun-flower, potato and fibre hemp; *vineyard and fruits*; and *other*: forest and lawn.

Area of agricultural land use shows clear, decreasing trend in the examined region (Fig. 2). In computation of area average evapotranspiration and albedo, however, variations of the total sown area are not considered, since the area-weighted sums will be normalised by this area, i.e. by the sum of the weights. On the other hand, possible variations at the set-aside areas are not included into our estimations.

4. Results and comparisons

4.1 Evapotranspiration

Change of land-use slightly increased evapotranspiration with strong fluctuations between 1951 and 2000 (Fig. 3.). Linear trend of the change was ca. 0.05 mm yr⁻¹, with an average increase of 2 mm for the total examined period (50 years). Between the two extremes of the curve there is a larger than 10 mm difference, but this is still not a dramatic change, since average evapotranspiration of the examined 7 months (Apr-Oct) is about 280 mm; that is to say, relative change is only 1 %.

On the basis of the above-mentioned Eq. (2), total fluctuation of evapotranspiration influenced by fluctuations of meteorological conditions was counted for the examined period between 1951 and 2000 (Mika et al., 2001). It could be seen, that inter-annual fluctuation of the total evapotranspiration exceeded effect of land use by one order of magnitude.

At the same time, there were no any one-way trends observed in the total evapotranspiration data sets. In summary of our results on evapotranspiration, it can be said that change of land use caused clear but little changes in the East-Hungarian region.

4.2 Surface albedo

Surface albedo multiplied with the yearly sown areas of each plant, give time series of albedo changes caused by changes in land use. The average surface albedo of the six counties shows decreasing tendency in each month from April to July (Mika et al., 2001).




Fig. 3. Effect of changes in land use to evapotranspiration (mm / 7 months).




Fig. 4. Effect of changes in land use to short wave energy reflected by the surface-atmosphere system, compared to the mean of the period 1951-80.

In other words, share of relatively darker plant stocks increased with larger green mass, giving thicker cover. The situation in August is the same, whereas no clear change can be established in the two following months. This may be in connection with the fact that plants, still present in September-October, do not develop any longer.

As it is assumed from the monthly decrease of albedo, regional averages of the surface energy balance increased steadily in the examined term. Trend of the change was $+0.017 \text{ Wm}^{-2}\text{yr}^{-1}$; namely, 0.75 Wm^{-2} as a total during the examined 50 years period. The linear trend fits fairly tightly to the data set, since value of the correlation coefficient is 0.98.

4.3 System albedo in comparison with other effects

If monthly values of the system albedo are multiplied by astronomically maximum sunshine radiation, coming to the outer boundary of the atmosphere, then short wave energy balance of the surface-atmosphere system is received. Energy surplus which remains in the system, because of the decrease of radiation energy reflected by the surface-atmosphere system (Fig. 4.), is $+0.010 \text{ Wm}^{-2}\text{yr}^{-1}$; namely, 0.5 Wm^{-2} as a total, during the 50 years.

In order to demonstrate the importance of this value, it should be noted that global atmospheric CO_2 -concentration has only shown 1.5 Wm^{-2} virtual increase per year since the industrial revolution. If the effect of concentration changes (from 310 to 357 ppm) is calculated by using a simplified zero-dimensional global model, parameterised from the experiments of recent (IPCC, 1996; p. 298, Table 6.3) GCMs, a smaller change is obtained: $+0.9 \text{ Wm}^{-2}$.

Change of albedo becomes more important, if it is considered that the same change of CO_2 -concentration, in relation to the average global long wave energy balance per year, has somewhat less effect in the summer in Hungary. The ratio, established by using the above radiative-convective model, is about 77 %; namely, the effect of change in CO_2 -concentration in Hungary during the examined 50 years period was only $+0.7 \text{ Wm}^{-2}$. Consequently, changes of land use between 1951 and 2000 caused more than half as much change in the energy balance of the surface-atmosphere system, than those of CO_2 -concentration.

It is worth calculating, how much is the effect of the 2 mm increase of evapotranspiration to heat release from the surface. Considering the fact, that evapotranspiration of 1 kg water needs 2,470 J energy, and the total 214 days of the examined seven months correspond to 18,489,600 seconds, this 2 mm evapotranspiration surplus deprives 0.25 Wm^{-2} energy of the energy balance of the surface. This value is of similar magnitude with that of change of CO_2 -concentration, detected during the examined 50 years period. Nevertheless, energy surplus used to evapotranspiration expresses only a minor contribution to modification of temperature at the different altitudes.

5. Discussion

The above calculations determined effect of changes in land use to evapotranspiration and albedo. These computations mixed fairly exact, *in situ* data on land use with generalised average physical parameters on albedo and biological constant of vegetation.

This might arise the question if the latter errors of estimation can influence the final conclusions, based on small differences of albedo and evapotranspiration. Although, it is rather difficult to quantify this effect, one may note that all annual values came from weighted averages of 20-25 different plants and that the parallel response of vegetation is plausible: the more heat income is compensated by enhanced evapotranspiration.

These effects of land use were compared to the components of the energy and water balance. According to these comparisons, it could be established that, for the region of East-Hungary ($34,000 \text{ km}^2$), changes due to land use were of the same magnitude with those of energy and water balance occurred in relation to the global climate change.

However, the question: whether or not changes in land use could produce that of regional climate (temperature, precipitation, etc.) with similar magnitude, could not be answered. To this, temporal change of land use should be known for a much larger region, as climate of Hungary is function of the energy- and water balance of a far larger area (this connection is less tight if the territory increases). That is to say, heat- and water balance strongly depend on horizontal transport of temperature and water vapour. Intensity of this procedure is, at first approach, directed by horizontal (spatial) differences of these two elements (temperature and water vapour). For this reason, if temporal change of the balances can be detected only at us, and not in the neighbouring regions, then this change is spread by the horizontal exchange on a well larger area.

In other words, the above result according to which changes of land use in East-Hungary in the last decades increased surface energy balance considerably but decreased water supply can be utilised if, together with similar examinations of the neighbouring countries, it might be built in a regional climate model.

Acknowledgements. The authors are indebted to L. Lukácsovics for his valuable contribution in series of computations; Z. Sümeghy for preparing Fig. 1.; A. Horváth for her contribution in the data collection and to I. Kevei-Bárány for professional advice on integrated classification of the wide variety of vegetation.

6. References

- Antal, E., 1968: *New method for the calculation of potential evaporation*. HMS Report on the Scientific Works in 1967. No.24, 414-423.
- Central Statistical Office, (1951-2000): *Statistical Yearbooks*. Central Statistical Office. County Centres of Hungary
- Charney, J.G., Quirk, W.J., Chow, S.H., and Kornfield, J., 1977: *A comparative study of the effects of albedo change on drought in semi-arid regions*. *J. Atmos. Sci.* 34, 1366-1385.
- CLIVAR, 1998: *Implementation Plan*. Hamburg, June, 1998, 260 p.
- Dávid, A., 1985: *Preliminary investigations estimating the regional distribution of surface albedo*. HMS Report on the Scientific Works in 1983. 81-98. (in Hungarian)
- Dunkel Z., 1994: *Investigation of climatic variability influence on soil moisture in Hungary*. In: XVIIth Conference of the Danube Countries, Budapest, Hungary, 441-446.

- Historical Statistical Contributions, (1971-79): *Data on Plant Management*. (Vol. I-IV.: 1922-33, 34-45, 47-62, 63-70.) Statisztikai Kiadó, Budapest. (in Hungarian)
- IPCC 1996: *Climate Change 1995. The Science of Climate Change*, the Second Assessment Report of the Intergovernmental Panel on Climate Change: Contribution of Working Group I. (Houghton, J. J., et al., eds.) Cambridge University Press, Cambridge.
- Karol I.L. and Frolkis A. A., 1984: *Energy-balance type radiative-convective model of the global climate*. Meteorology and Hydrology, No. 8, 59-68. (in Russian)
- Mika, J., 1988, *Regional features of a global warming in the Carpathian Basin*. Időjárás 92, 178-189. (in Hungarian)
- Mika J., Kovács E., Németh P., and Rimóczi-Paál A., 1991: *Parameterisation for regional energy balance climate modeling over Hungary*. Adv. in Space Res. 11, (3)101-(3)104.
- Mika J., Kovács E., and Dávid A., 1992: *Effect of vegetation and soil conditions on albedo of the surface-atmosphere system*. HMS Report on the Scientific Works in 1988, 165-173. (in Hungarian)
- Mika J., Kovács E., and Bozó L., 1993: *Effects of anthropogenic and natural changes in atmospheric aerosol content on radiation balance in the Carpathian Basin*. HMS Report on the Scientific Works in 1989, 81-89. (in Hungarian)
- Mika J., Horváth, Sz., Fogarasi J. and Makra, L., 1998: *Simulation of climate forcing mechanisms on the energy and water balance of a watershed*. In: 19th Conf. Danube Countries, Osijek, Croatia, 15-19, June 1998, 331-342.
- Mika J., Horváth Sz. and Makra L., 2001: *Impact of documented land-use changes on the surface albedo and evapotranspiration in a plain watershed*. Phys. Chem. Earth (B), Vol. 26, No. 7-8, 601-606.
- Posza, I. and Stollár, A. 1983: Plant constants evaluated by several years' measurements for the calculation of effective evaporation. Időjárás 87, 170-177. (in Hungarian)
- Práger T. and Kovács E., 1988: *Investigation of effects of atmospheric trace gases and aerosol on climate with a radiative-convective model*. Időjárás 92 153-162. (in Hungarian)
- Sagan, C., Toon, O.B., and Pollack, J.B., 1979: *Anthropogenic albedo changes and the earth's climate*. Science, 206, 1363-68.
- Warren, S.G., Hahn, C.J., and London, J., 1985: *Simultaneous Occurrence of Different Cloud Types*. J. Climate Appl. Met. 24, 658-667.
- WMO, 1983: *Report of the Experts Meeting on Aerosols and Their Climatic Effects*. Williamsburg, Virginia, 28-30 March 1983, World Climate Programme, WCP-55.